

Scale Invariance and Vacuum Energy

E. I. Guendelman¹

Physics Department, Ben Gurion University, Beer Sheva, Israel

Abstract

The possibility of mass in the context of scale-invariant, generally covariant theories, is discussed. Scale invariance is considered in the context of a gravitational theory where the action, in the first order formalism, is of the form $S = \int L_1 \Phi d^4x + \int L_2 \sqrt{-g} d^4x$ where Φ is a density built out of degrees of freedom independent of the metric. For global scale invariance, a "dilaton" ϕ has to be introduced, with non-trivial potentials $V(\phi) = f_1 e^{\alpha\phi}$ in L_1 and $U(\phi) = f_2 e^{2\alpha\phi}$ in L_2 . This leads to non-trivial mass generation and a potential for ϕ which is interesting for new inflation. Scale invariant mass terms for fermions lead to a possible explanation of the present day accelerated universe and of cosmic coincidences.

The concept of scale invariance appears as an attractive possibility for a fundamental symmetry of nature. In its most naive realizations, such a symmetry is not a viable symmetry, however, since nature seems to have chosen some typical scales.

Here we will find that scale invariance can nevertheless be incorporated into realistic, generally covariant field theories. However, scale invariance has to be discussed in a more general framework than that of standard generally relativistic theories, where we must allow in the action, in addition to the ordinary measure of integration $\sqrt{-g} d^4x$, another one, Φd^4x , where Φ is a density built out of degrees of freedom independent of the metric.

For example, given 4-scalars φ_a ($a = 1, 2, 3, 4$), one can construct the density

$$\Phi = \varepsilon^{\mu\nu\alpha\beta} \varepsilon_{abcd} \partial_\mu \varphi_a \partial_\nu \varphi_b \partial_\alpha \varphi_c \partial_\beta \varphi_d \quad (1)$$

¹Electronic address: guendel@bgumail.bgu.ac.il

One can allow both geometrical objects to enter the theory and consider¹

$$S = \int L_1 \Phi d^4x + \int L_2 \sqrt{-g} d^4x \quad (2)$$

Here L_1 and L_2 are φ_a independent. There is a good reason not to consider mixing of Φ and $\sqrt{-g}$, like for example using $\frac{\Phi^2}{\sqrt{-g}}$. This is because (2) is invariant (up to the integral of a total divergence) under the infinite dimensional symmetry $\varphi_a \rightarrow \varphi_a + f_a(L_1)$ where $f_a(L_1)$ is an arbitrary function of L_1 if L_1 and L_2 are φ_a independent. Such symmetry (up to the integral of a total divergence) is absent if mixed terms are present.

We will study now the dynamics of a scalar field ϕ interacting with gravity as given by the action (2) with²

$$L_1 = \frac{-1}{\kappa} R(\Gamma, g) + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi), L_2 = U(\phi) \quad (3)$$

$$R(\Gamma, g) = g^{\mu\nu} R_{\mu\nu}(\Gamma), R_{\mu\nu}(\Gamma) = R_{\mu\nu\lambda}^\lambda(\Gamma) = \Gamma_{\mu\nu,\sigma}^\lambda - \Gamma_{\mu\sigma,\nu}^\lambda + \Gamma_{\alpha\sigma}^\lambda \Gamma_{\mu\nu}^\alpha - \Gamma_{\alpha\nu}^\lambda \Gamma_{\mu\sigma}^\alpha. \quad (4)$$

In the variational principle $\Gamma_{\mu\nu}^\lambda, g_{\mu\nu}$, the measure fields scalars φ_a and the scalar field ϕ are all to be treated as independent variables.

If we perform the global scale transformation ($\theta = \text{constant}$)

$$g_{\mu\nu} \rightarrow e^\theta g_{\mu\nu} \quad (5)$$

then (2), with the definitions (3), (4), is invariant provided $V(\phi)$ and $U(\phi)$ are of the form

$$V(\phi) = f_1 e^{\alpha\phi}, U(\phi) = f_2 e^{2\alpha\phi} \quad (6)$$

and φ_a is transformed according to $\varphi_a \rightarrow \lambda_a \varphi_a$ (no sum on a) which means $\Phi \rightarrow \left(\prod_a \lambda_a\right) \Phi \equiv \lambda \Phi$ such that $\lambda = e^\theta$ and $\phi \rightarrow \phi - \frac{\theta}{\alpha}$. In this case we call the scalar field ϕ needed to implement scale invariance "dilaton".

Let us consider the equations which are obtained from the variation of the φ_a fields. We obtain then $A_a^\mu \partial_\mu L_1 = 0$ where $A_a^\mu = \varepsilon^{\mu\nu\alpha\beta} \varepsilon_{abcd} \partial_\nu \varphi_b \partial_\alpha \varphi_c \partial_\beta \varphi_d$. Since $\det(A_a^\mu) = \frac{4!}{4!} \Phi^3 \neq 0$ if $\Phi \neq 0$. Therefore if $\Phi \neq 0$ we obtain that $\partial_\mu L_1 = 0$, or that $L_1 = M$, where M is constant. This constant M appears in a self-consistency condition of the equations of motion that allows us to solve for $\chi \equiv \frac{\Phi}{\sqrt{-g}}$

$$\chi = \frac{2U(\phi)}{M + V(\phi)}. \quad (7)$$

To get the physical content of the theory, it is convenient to go to the Einstein conformal frame where

$$\bar{g}_{\mu\nu} = \chi g_{\mu\nu} \quad (8)$$

and χ given by (7). In terms of $\bar{g}_{\mu\nu}$ the non Riemannian contribution (defined as $\Sigma_{\mu\nu}^\lambda = \Gamma_{\mu\nu}^\lambda - \{\lambda_{\mu\nu}\}$ where $\{\lambda_{\mu\nu}\}$ is the Christoffel symbol), disappears from the equations, which can be written then in the Einstein form ($R_{\mu\nu}(\bar{g}_{\alpha\beta}) =$ usual Ricci tensor)

$$R_{\mu\nu}(\bar{g}_{\alpha\beta}) - \frac{1}{2}\bar{g}_{\mu\nu}R(\bar{g}_{\alpha\beta}) = \frac{\kappa}{2}T_{\mu\nu}^{eff}(\phi) \quad (9)$$

where

$$T_{\mu\nu}^{eff}(\phi) = \phi_{,\mu}\phi_{,\nu} - \frac{1}{2}\bar{g}_{\mu\nu}\phi_{,\alpha}\phi_{,\beta}\bar{g}^{\alpha\beta} + \bar{g}_{\mu\nu}V_{eff}(\phi), V_{eff}(\phi) = \frac{1}{4U(\phi)}(V + M)^2. \quad (10)$$

If $V(\phi) = f_1 e^{\alpha\phi}$ and $U(\phi) = f_2 e^{2\alpha\phi}$ as required by scale invariance, we obtain from (10)

$$V_{eff} = \frac{1}{4f_2}(f_1 + M e^{-\alpha\phi})^2 \quad (11)$$

Since we can always perform the transformation $\phi \rightarrow -\phi$ we can choose by convention $\alpha > 0$. We then see that as $\phi \rightarrow \infty$, $V_{eff} \rightarrow \frac{f_1^2}{4f_2} = \text{const.}$ providing an infinite flat region. Also a minimum is achieved at zero cosmological constant for the case $\frac{f_1}{M} < 0$ at the point $\phi_{min} = \frac{-1}{\alpha} \ln \left| \frac{f_1}{M} \right|$. Finally, the second derivative of the potential V_{eff} at the minimum is $V_{eff}'' = \frac{\alpha^2}{2f_2} |f_1|^2 > 0$ if $f_2 > 0$,

There are many interesting issues that one can raise here. The first one is of course the fact that a realistic scalar field potential, with massive excitations when considering the true vacuum state, is achieved in a way consistent with the idea of scale invariance.

The second point to be raised is that since there is an infinite region of flat potential for $\phi \rightarrow \infty$, we expect a slow rolling new inflationary³ scenario to be viable, provided the universe is started at a sufficiently large value of the scalar field ϕ .

Furthermore, one can consider this model as suitable for the present day universe rather than for the early universe, after we suitably reinterpret the meaning of the scalar field ϕ . This can provide a long lived almost constant vacuum energy for a long period of time, which can be small if $f_1^2/4f_2$ is small. Such small energy density will eventually disappear when the universe achieves its true vacuum state.

Notice that a small value of $\frac{f_1^2}{f_2}$ can be achieved if we let $f_2 \gg f_1$. In this case $\frac{f_1^2}{f_2} \ll f_1$, i.e. a very small scale for the energy density of the universe is obtained by the existence of a very high scale (that of f_2) the same way as a small fermion mass is obtained in the see-saw mechanism⁴ from the existence also of a large mass scale. In what follows, we will take $f_2 \gg f_1$.

So far we have studied a theory which contains the metric tensor $g_{\mu\nu}$, the measure fields φ_a (a=1,2,3,4) and the "dilaton" ϕ , which makes global scale invariance possible in a non-trivial way. All of the above fields have some kind of geometrical significance, but if we are to describe the real world, the list of fields and/or particles has to be enlarged.

Taking, for example, the case of a fermion ψ , where the kinetic term of the fermion is chosen to be part of L_1

$$S_{fk} = \int L_{fk} \Phi d^4x \quad (12)$$

$$L_{fk} = \frac{i}{2} \bar{\psi} [\gamma^a V_a^\mu (\vec{\partial}_\mu + \frac{1}{2} \omega_\mu^{cd} \sigma_{cd}) - (\overleftarrow{\partial}_\mu + \frac{1}{2} \omega_\mu^{cd} \sigma_{cd}) \gamma^a V_a^\mu] \psi \quad (13)$$

there V_a^μ is the vierbein, $\sigma_{cd} = \frac{1}{2} [\gamma_c, \gamma_d]$, the spin connection ω_μ^{cd} is determined by variation with respect to ω_μ^{cd} and, for self-consistency, the curvature scalar is taken to be (if we want to deal with ω_μ^{ab} instead of $\Gamma_{\mu\nu}^\lambda$ everywhere)

$$R = V^{a\mu} V^{b\nu} R_{\mu\nu ab}(\omega), R_{\mu\nu ab}(\omega) = \partial_\mu \omega_{\nu ab} - \partial_\nu \omega_{\mu ab} + (\omega_{\mu a}^c \omega_{\nu cb} - \omega_{\nu a}^c \omega_{\mu cb}). \quad (14)$$

Global scale invariance is obtained provided ψ also transforms, as in $\psi \rightarrow \lambda^{-\frac{1}{4}} \psi$. Mass term consistent with scale invariance exist,

$$S_{fm} = m_1 \int \bar{\psi} \psi e^{\alpha\phi/2} \Phi d^4x + m_2 \int \bar{\psi} \psi e^{3\alpha\phi/2} \sqrt{-g} d^4x. \quad (15)$$

If we consider the situation where $m_1 e^{\alpha\phi/2} \bar{\psi} \psi$ or $m_2 e^{3\alpha\phi/2} \bar{\psi} \psi$ are much bigger than $V(\phi) + M$, i.e. a high density approximation, we obtain that

instead of (7) that the consistency condition is¹ $(3m_2e^{3\alpha\phi/2} + m_1e^{\alpha\phi/2}\chi)\bar{\psi}\psi = 0$, which means $\chi = -\frac{3m_2}{m_1}e^{\alpha\phi}$. Using this in (15), we obtain, after going to the conformal Einstein frame, which involves, also a transformation of the fermion fields, necessary so as to achieve Einstein-Cartan form for both the gravitational and fermion equations. These transformations are, $\bar{g}_{\mu\nu} = \chi g_{\mu\nu}$ (or $\bar{V}_\mu^a = \chi^{\frac{1}{2}}V_\mu^a$) and $\psi' = \chi^{-\frac{1}{4}}\psi$ and they lead to a mass term,

$$S_{fm} = -2m_2\left(\frac{|m_1|}{3|m_2|}\right)^{3/2} \int \sqrt{-\bar{g}}\bar{\psi}'\psi' d^4x \quad (16)$$

The ϕ dependence of the mass term has disappeared, i.e. masses are constants.

There is one situation where the low density of matter can also give results which are similar to those obtained in the high density approximation, in that the coupling of the ϕ field disappears and that the mass term becomes of a conventional form in the Einstein conformal frame.

This is the case, when we study the theory for the limit $\phi \rightarrow \infty$. Then $U(\phi) \rightarrow \infty$ and $V(\phi) \rightarrow \infty$. In this case, taking $m_1e^{\alpha\phi/2}\bar{\psi}\psi$ and $m_2e^{3\alpha\phi/2}\bar{\psi}\psi$ much smaller than $V(\phi)$ or $U(\phi)$ respectively, therefore one can see that (7) is a good approximation and since also M can be ignored in the self consistency condition (7) in this limit, we get then, $\chi = \frac{2f_2}{f_1}e^{\alpha\phi}$. If this is inserted in (15), we get $S_{fm} = m \int \sqrt{-\bar{g}}\bar{\psi}'\psi' d^4x$, where

$$m = m_1\left(\frac{f_1}{2f_2}\right)^{\frac{1}{2}} + m_2\left(\frac{f_1}{2f_2}\right)^{\frac{3}{2}} \quad (17)$$

Comparing (16) and (17) and taking m_1 and m_2 of the same order of magnitude, we see that the mass of the Dirac particle is much smaller in the region $\phi \rightarrow \infty$, for which (17) is valid, than it is in the region of high density of the Dirac particle relative to $V(\phi) + M$, as displayed in eq. (16), if the "see-saw" assumption $\frac{f_1}{f_2} \ll 1$ is made.

Therefore if space is populated by these diluted Dirac particles of this type, the mass of these particles will grow substantially if we go to the true vacuum state valid in the absence of matter, i.e. $V + M = 0$, as dictated by V_{eff} given by eq. (11). The presence of matter pushes therefore the minimum of energy to a state where $V + M > 0$. The real vacuum in the presence of matter should not be located in the region $\phi \rightarrow \infty$, which minimizes the matter energy, but maximizes the potential energy V_{eff} and

not at $V + M = 0$, which minimizes V_{eff} , and where particle masses are big, but somewhere in a balanced intermediate stage. Clearly how much above $V + M = 0$ such true vacuum is located must be correlated to how much particle density is there in the Universe. A non zero vacuum energy, which must be of the same order of the particle energy density, has to appear and this could explain the "accelerated universe" that appears to be implied by the most recent observations, together with the "cosmic coincidence", that requires the vacuum energy be of the same order of magnitude to the matter energy⁵.

I would like to thank J. Bekenstein, A. Davidson, A.Guth and A. Kaganovich for conversations on the subjects discussed here.

References

- [1] For a review and further references to published papers (in Phys. Rev. D., Mod. Phys. Lett.A, etc.) see, E.I.Guendelman and A.B.Kaganovich, Phys.Rev.D.60, 065004 (1999).
- [2] E.I. Guendelman, gr-qc/9901067; E.I. Guendelman, Mod. Phys. Lett. A14: 1043 (1999) (gr-qc/9901017).
- [3] A.D. Linde, Phys. Lett, 108B, (1982) 389; A. Albrecht and P.J. Steinhardt, Phys. Rev. Lett, 48, (1982) 1220.
- [4] For a review see B.Kayser, The physics of massive neutrinos, World Scientific, Singapore (1989).
- [5] For a review see articles by C.J.Hogan, R.P.Krishner and N.B.Suntzeff in Sci. Amer. 280 (1999), 28 and by L.Krauss, Sci. Amer. 280, (1999) 34.